

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C11D 1/65, 1/86	A1	(11) International Publication Number: WO 97/12018 (43) International Publication Date: 3 April 1997 (03.04.97)
(21) International Application Number: PCT/US95/12470 (22) International Filing Date: 29 September 1995 (29.09.95) (71) Applicant (for all designated States except US): THE PROCTER & GAMBLE COMPANY [US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): LAPPAS, Dimitris [GR/BE]; 5, rue Alphons Wauters, B-1020 Brussels (BE). PANANDIKER, Rajan, Keshav [IN/US]; 1321 Williamsburg Road, Cincinnati, OH 45215 (US). HORNER, Thomas, Wilhelm [DE/BE]; Hindelaan 3A, B-3090 Overijse (BE). (74) Agents: REED, T., David et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45217 (US).		(81) Designated States: AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, IS, JP, KG, KP, KR, KZ, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TT, UA, US, UZ, VN, ARIPO patent (KE, MW, SD, SZ, UG), European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: LIQUID LAUNDRY DETERGENTS CONTAINING SELECTED QUATERNARY AMMONIUM COMPOUNDS (57) Abstract <p>This invention relates to a liquid laundry detergent composition comprising a surfactant system which is free of linear alkyl benzene sulfonate comprising: i) anionic surfactants selected from the group of alkyl alkoxy sulfates and alkyl sulfates; ii) a selected quaternary ammonium surfactant.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

LIQUID LAUNDRY DETERGENTS
CONTAINING
SELECTED QUATERNARY AMMONIUM COMPOUNDS

Field of the Invention

The present invention relates to detergent compositions containing surfactants selected from quaternary ammonium surfactants.

More particularly, the invention is directed to detergent compositions containing a nil-LAS surfactant system comprising anionic surfactants selected from the group consisting of alkyl alkoxylated sulfates and alkyl sulfates, said composition further containing specific quaternary ammonium surfactants.

Background of the Invention

The present invention relates to liquid detergents containing a surfactant system which is free of Linear alkyl benzene sulfonate surfactants (LAS), said liquid detergent having optimum greasy stain removal performance.

The recent trend towards partial or total replacement of Linear alkyl benzene sulfonate surfactants (LAS) has

urged the detergent formulators to rebalance their formulations with different surfactants.

There is thus a standing desire for performance and flexibility reasons to make available a surfactant system capable of providing optimum detergency performance which is equivalent to that of LAS-containing detergents.

The above objective has been met by a surfactant system comprising anionic surfactants selected from the group consisting of alkyl alkoxylated sulfates and alkyl sulfates, said surfactant system further comprising a cosurfactant selected from the group of quaternary ammonium surfactants.

It has been surprisingly found that detergent compositions containing said surfactant system exhibit detergency performance equivalent to that of LAS-containing detergents.

In addition, it was found that the liquid detergent compositions containing the selected quaternary ammonium surfactants of the present invention, provide excellent greasy stain removal performance without detriment to the suds characteristics of the compositions. This finding allows to reduce the level of suds suppressing agents, thereby facilitating the formulation of concentrated liquid detergents.

Quaternary ammonium surfactants are described in the art. The properties of these surfactants are very strongly influenced by the type of substituent. Chain length, degree of saturation, branching or the presence and number of hydroxylic or ethoxy groups mainly determine the properties of the surfactant. Whereas typical textile-conditioning actions are performed by cationic surfactants with two long alkyl chains, cationic surfactants with only one long alkyl chain have been reported to improve the detergency performance in laundry detergents. EP-A-224 describes liquid built laundry detergent compositions comprising a general class of quaternary ammonium surfactants. Decyltrimethyl

ammonium chloride is described. EP 8142 describes a liquid builder-free heavy duty detergent comprising a quaternary ammonium compound of a general formula. Octyldihydroxyethylmethyl ammonium halides are described.

For optimum grease detergency performance, however, these compositions of the prior art require high level of cationic surfactant. These high levels of cationic surfactants in turn, generate excessive foaming, thereby raising problems of automatic washing machine compatibility. If, on the other hand, the cationic surfactant is reduced at a level at which foam regulation is no longer a problem, the beneficial grease detergency characteristics of quaternary compounds are diminished.

In contrast, the surfactant system of the present invention provides optimum grease and oil removal performance, thereby not adversely affecting the suds characteristic of the detergent compositions formulated therewith.

In addition, it has been found that liquid detergent compositions formulated with said surfactant system are extremely useful when the liquid detergent compositions are in direct contact with the fabrics such as during pretreatment.

Summary of the Invention

The present invention relates to liquid detergent compositions comprising a Nil-LAS surfactant system said surfactant system comprising anionic surfactants selected from the group of alkyl alkoxy sulfates and alkyl sulfates and selected quaternary ammonium surfactants present in specific weight ratio.

The detergent compositions preferably comprise at least 5%, more preferably from 10% to 65% and most preferably from 15% to 40% by weight of the surfactant system as described hereinabove.

Detailed description of the Invention

The surfactant system of the detergent compositions according to the present invention comprise anionic surfactants selected from the group of alkylalkoxy sulfates and alkyl sulfates.

Alkyl alkoxyated sulfates and/or alkyl sulfates


The alkyl alkoxyated sulfate surfactants hereof are water soluble salts or acids of the formula $RO(A)_mSO_3M$ wherein R is an unsubstituted C_{10} - C_{24} alkyl or hydroxyalkyl group having a C_{10} - C_{24} alkyl component, preferably a C_{12} - C_{18} alkyl or hydroxyalkyl, more preferably C_{12} - C_{15} alkyl or hydroxyalkyl, A is an ethoxy or propoxy unit, m is greater than zero, typically between about 0.5 and about 6, more preferably between about 0.5 and about 3, and M is H or a cation which can be, for example, a metal cation (e.g., sodium, potassium, lithium, calcium, magnesium, etc.), ammonium or substituted-ammonium cation. Alkyl ethoxyated sulfates as well as alkyl propoxyated sulfates are contemplated herein. Specific examples of substituted ammonium cations include ethanol-, triethanol-, methyl-, dimethyl, trimethyl-ammonium cations and quaternary ammonium cations such as tetramethyl-ammonium and dimethyl piperidinium cations and those derived from alkylamines such as ethylamine, diethylamine, triethylamine, mixtures thereof, and the like. Exemplary surfactants are C_{12} - C_{15} alkyl polyethoxylate (1.0) sulfate (C_{12} - $C_{15}E(1.0)M$), C_{12} - C_{15} alkyl polyethoxylate (2.25) sulfate (C_{12} - $C_{15}E(2.25)M$), C_{12} -

C₁₅ alkyl polyethoxylate (3.0) sulfate (C₁₂-C₁₅E(3.0)M), and C₁₂-C₁₅ alkyl polyethoxylate (4.0) sulfate (C₁₂-C₁₅E(4.0)M), wherein M is conveniently selected from sodium and potassium.

The alkyl sulfate surfactants hereof are water soluble salts or acids of the formula ROSO₃M wherein R preferably is a C₁₀-C₂₄ hydrocarbyl, preferably an alkyl or hydroxyalkyl having a C₁₀-C₁₈ alkyl component, more preferably a C₁₂-C₁₅ alkyl or hydroxyalkyl, and M is H or a cation, e.g., an alkali metal cation (e.g. sodium, potassium, lithium), or ammonium or substituted ammonium (e.g. methyl-, dimethyl-, and trimethyl ammonium cations and quaternary ammonium cations such as tetramethyl-ammonium and dimethyl piperidinium cations and quaternary ammonium cations derived from alkylamines such as ethylamine, diethylamine, triethylamine, and mixtures thereof, and the like).


The quaternary ammonium compound

The quaternary ammonium surfactant according to the present invention has the formula (I):



Formula I

whereby R₁ is a short chainlength alkyl (C₆-C₁₀) or alkylamidoalkyl of the formula (II) :



Formula II


y is 2-4, preferably 3.

whereby R₂ is H or a C₁-C₃ alkyl,

whereby x is 0-4, preferably 0-2, most preferably 0,

whereby R₃, R₄ and R₅ are either the same or different and can be either a short chain alkyl (C₁-C₃) or alkoxyated alkyl of the formula III,

whereby X⁻ is a counterion, preferably a halide, e.g. chloride or methylsulfate.



Formula III

R₆ is C₁-C₄ and z is 1 or 2.

Preferred quat ammonium surfactants are those as defined in formula I whereby

R₁ is C₈, C₁₀ or mixtures thereof, x=0,

R₃, R₄ = CH₃ and R₅ = CH₂CH₂OH.

Preferred quaternary ammonium surfactants have the general formula I, whereby the weight ratio of the quaternary ammonium surfactant to the alkyl alkoxy sulfates and alkyl sulfates is from 1:3 to 1:30, preferably from 1:3 to 1:15, most preferred from 1:5 to 1:10.

Detergent ingredients

In another embodiment of the present invention, a liquid detergent composition is provided comprising the surfactant system of the present invention mixed with detergent ingredients. A wide range of surfactants can be used in the detergent composition of the present invention. The detergent compositions according to the present invention comprise a surfactant system which is

substantially free of linear alkylbenzene sulfonate surfactant.

A typical listing of anionic, nonionic, ampholytic and zwitterionic classes, and species of these surfactants, is given in US Patent 3,664,961 issued to Norris on May 23, 1972.

Other suitable anionic surfactants that can be used are alkyl ester sulfonate surfactants including linear esters of C₈-C₂₀ carboxylic acids (i.e., fatty acids) which are sulfonated with gaseous SO₃ according to "The Journal of the American Oil Chemists Society", 52 (1975), pp. 323-329. Suitable starting materials would include natural fatty substances as derived from tallow, palm oil, etc.

The preferred alkyl ester sulfonate surfactant, especially for laundry applications, comprise alkyl ester sulfonate surfactants of the structural formula :



wherein R³ is a C₈-C₂₀ hydrocarbyl, preferably an alkyl, or combination thereof, R⁴ is a C₁-C₆ hydrocarbyl, preferably an alkyl, or combination thereof, and M is a cation which forms a water soluble salt with the alkyl ester sulfonate. Suitable salt-forming cations include metals such as sodium, potassium, and lithium, and substituted or unsubstituted ammonium cations, such as monoethanolamine, diethanolamine, and triethanolamine. Preferably, R³ is C₁₀-C₁₆ alkyl, and R⁴ is methyl, ethyl or isopropyl. Especially preferred are the methyl ester sulfonates wherein R³ is C₁₀-C₁₆ alkyl.

Other anionic surfactants useful for deterative purposes can also be included in the laundry detergent compositions of the present invention. These can include salts (including, for example, sodium, potassium, ammonium, and substituted ammonium salts such as mono-, di- and


triethanolamine salts) of soap, C₈-C₂₂ primary or secondary alkanesulfonates, C₈-C₂₄ olefinsulfonates, sulfonated polycarboxylic acids prepared by sulfonation of the pyrolyzed product of alkaline earth metal citrates, e.g., as described in British patent specification No. 1,082,179, C₈-C₂₄ alkylpolyglycoethersulfates (containing up to 10 moles of ethylene oxide); alkyl glycerol sulfonates, fatty acyl glycerol sulfonates, fatty oleoyl glycerol sulfates, alkyl phenol ethylene oxide ether sulfates, paraffin sulfonates, alkyl phosphates, isethionates such as the acyl isethionates, N-acyl taurates, alkyl succinamates and sulfosuccinates, monoesters of sulfosuccinates (especially saturated and unsaturated C₁₂-C₁₈ monoesters) and diesters of sulfosuccinates (especially saturated and unsaturated C₆-C₁₂ diesters), sulfates of alkylpolysaccharides such as the sulfates of alkylpolyglucoside (the nonionic nonsulfated compounds being described below), and alkyl polyethoxy carboxylates such as those of the formula $RO(CH_2CH_2O)_k-CH_2COO-M^+$ wherein R is a C₈-C₂₂ alkyl, k is an integer from 0 to 10, and M is a soluble salt-forming cation. Resin acids and hydrogenated resin acids are also suitable, such as rosin, hydrogenated rosin, and resin acids and hydrogenated resin acids present in or derived from tall oil. Further examples are described in "Surface Active Agents and Detergents" (Vol. I and II by Schwartz, Perry and Berch). A variety of such surfactants are also generally disclosed in U.S. Patent 3,929,678, issued December 30, 1975 to Laughlin, et al. at Column 23, line 58 through Column 29, line 23 (herein incorporated by reference).

When included therein, the laundry detergent compositions of the present invention typically comprise from about 1% to about 40%, preferably from about 5% to about 25% by weight of such anionic surfactants.

One class of nonionic surfactants useful in the present invention are condensates of ethylene oxide with a hydrophobic moiety to provide a surfactant having an average hydrophilic-lipophilic balance (HLB) in the range from 8 to 17, preferably from 9.5 to 14, more preferably from 12 to 14. The hydrophobic (lipophilic) moiety may be aliphatic or aromatic in nature and the length of the polyoxyethylene group which is condensed with any particular hydrophobic group can be readily adjusted to yield a water-soluble compound having the desired degree of balance between hydrophilic and hydrophobic elements.

Especially preferred nonionic surfactants of this type are the C₉-C₁₅ primary alcohol ethoxylates containing 3-12 moles of ethylene oxide per mole of alcohol, particularly the C₁₂-C₁₅ primary alcohols containing 5-8 moles of ethylene oxide per mole of alcohol.

Another class of nonionic surfactants comprises alkyl polyglucoside compounds of general formula




wherein Z is a moiety derived from glucose; R is a saturated hydrophobic alkyl group that contains from 12 to 18 carbon atoms; t is from 0 to 10 and n is 2 or 3; x is from 1.3 to 4, the compounds including less than 10% unreacted fatty alcohol and less than 50% short chain alkyl polyglucosides. Compounds of this type and their use in detergent are disclosed in EP-B 0 070 077, 0 075 996 and 0 094 118.

Very suitable as nonionic surfactants are poly hydroxy fatty acid amide surfactants of the formula



wherein R^1 is H, or R^1 is C_{1-4} hydrocarbyl, 2-hydroxy ethyl, 2-hydroxy propyl or a mixture thereof, R^2 is C_{5-31} hydrocarbyl, and Z is a polyhydroxyhydrocarbyl having a linear hydrocarbyl chain with at least 3 hydroxyls directly connected to the chain, or an alkoxyated derivative thereof. Preferably, R^1 is methyl, R^2 is a straight C_{11-15} alkyl or alkenyl chain such as coconut alkyl or mixtures thereof, and Z is derived from a reducing sugar such as glucose, fructose, maltose, lactose, in a reductive amination reaction.

Highly preferred nonionics are amine oxide surfactants. The compositions of the present invention may comprise amine oxide in accordance with the general formula I:



In general, it can be seen that the structure (I) provides one long-chain moiety $R^1 (EO)_x (PO)_y (BO)_z$ and two short chain moieties, $CH_2 R'$. R' is preferably selected from hydrogen, methyl and $-CH_2OH$. In general R^1 is a primary or branched hydrocarbyl moiety which can be saturated or unsaturated, preferably, R^1 is a primary alkyl moiety. When $x+y+z = 0$, R^1 is a hydrocarbyl moiety having chainlength of from about 8 to about 18. When $x+y+z$ is different from 0, R^1 may be somewhat longer, having a chainlength in the range $C_{12}-C_{24}$. The general formula also encompasses amine oxides wherein $x+y+z = 0$, $R^1 = C_8-C_{18}$, $R' = H$ and $q = 0-2$, preferably 2. These amine oxides are illustrated by $C_{12}-14$ alkyldimethyl amine oxide, hexadecyl dimethylamine oxide, octadecylamine oxide and their hydrates, especially the dihydrates as disclosed in U.S. Patents 5,075,501 and 5,071,594, incorporated herein by reference.

The invention also encompasses amine oxides wherein $x+y+z$ is different from zero, specifically $x+y+z$ is from

about 1 to about 10, R^1 is a primary alkyl group containing 8 to about 24 carbons, preferably from about 12 to about 16 carbon atoms; in these embodiments $y + z$ is preferably 0 and x is preferably from about 1 to about 6, more preferably from about 2 to about 4; EO represents ethyleneoxy; PO represents propyleneoxy; and BO represents butyleneoxy. Such amine oxides can be prepared by conventional synthetic methods, e.g., by the reaction of alkylethoxysulfates with dimethylamine followed by oxidation of the ethoxylated amine with hydrogen peroxide.

Highly preferred amine oxides herein are solids at ambient temperature, more preferably they have melting-points in the range 30°C to 90°C. Amine oxides suitable for use herein are made commercially by a number of suppliers, including Akzo Chemie, Ethyl Corp., and Procter & Gamble. See McCutcheon's compilation and Kirk-Othmer review article for alternate amine oxide manufacturers. Preferred commercially available amine oxides are the solid, dihydrate ADMOX 16 and ADMOX 18, ADMOX 12 and especially ADMOX 14 from Ethyl Corp.


Preferred embodiments include hexadecyldimethylamine oxide dihydrate, dodecyldimethylamine oxide dihydrate, octadecyldimethylamine oxide dihydrate, hexadecyltris(ethyleneoxy)dimethyl-amine oxide, and tetradecyldimethyl-amine oxide dihydrate.

Whereas in certain of the preferred embodiments $R' = H$, there is some latitude with respect to having R' slightly larger than H. Specifically, the invention further encompasses embodiments wherein $R' = CH_2OH$, such as hexadecylbis(2-hydroxyethyl)amine oxide, ²tallowbis(2-hydroxyethyl)amine oxide, stearylbis(2-hydroxyethyl)amine oxide and oleylbis(2-hydroxyethyl)amine oxide, dodecyldimethylamine oxide dihydrate.

Suitable nonionic surfactants include primary amines according to the formula $R_1R_2R_3N$ wherein R_1 and R_2 are both H, R_3 is a C_4 - C_{18} , preferably C_6 - C_{12} alkyl chain, R_3 alkyl chains may be straight or branched and may be interrupted with up to 12 ethylene oxide moieties, most preferably interrupted with up to 5 ethylene oxide moieties. Preferred amines according to the formula herein above are n-alkyl amines. Suitable amines for use herein may be selected from 1-hexylamine, 1-octylamine, laurylamine, palmitylamine, stearylamine, oleylamine, coconutalkylamine, tallowalkylamine.

Other suitable primary amines include amines according to the formula $R_1R_2R_3N$ wherein R_1 and R_2 are both H ; R_3 is $R_4X(CH_2)_n$, X is $-O-$, $-C(O)NH-$ or $-NH-$, R_4 is a C_4 - C_{18} , preferably C_6 - C_{12} alkyl chain and R_4 may be branched or straight, n is between 1 to 5. Preferred amines according to the formula herein above 3-isopropoxypropylamine, 3-(2-methoxyethoxy)-propylamine and 2-(2-aminoethoxy)-ethanol, C_8 - C_{10} octyl oxy propylamine, 2-ethylhexyloxypropylamine, lauryl amido propylamine and coco amido propylamine.

Suitable tertiary amines for use herein include amines tertiary amines having the formula $R_1R_2R_3N$ wherein neither R_1 nor R_2 is H, R_1 and R_2 are C_1 - C_8 alkylchains or



whereby n is between 2 to 4 and x is between 1 to 6;


R_3 is either a C_4 - C_{18} , preferably C_6 - C_{12} alkyl chain, or R_3 is $R_4X(CH_2)_n$, whereby X is $-O-$, $-C(O)NH-$ or $-NH-$, R_4 is a C_4 - C_{18} , n is between 1 to 5, and R_5 is H or C_1 - C_2 alkyl.

R_3 , R_4 are preferably C_6 - C_{12} alkyl chains and may be straight or branched ; R_3 alkyl chains may be interrupted with up to 12 ethylene oxide moieties, most preferably interrupted with up to 5 ethylene oxide moieties.

Suitable tertiary amines for use herein include coconutalkyldimethylamine, dimethyloleylamine, hexadecyltris (ethyleneoxy)dimethylamine, tallowalkylbis(2-hydroxyethyl)amine, stearylbis(2-hydroxyethyl)amine and oleoylbis(2-hydroxyethyl)amine.

Of all of the foregoing amines the preferred materials are the trialkyl amines marketed under the tradename ADOGEN, the long chain alkyldimethyl amines marketed under the tradename ARMEEN and the ethoxylated amines marketed under the tradename ETHOMEEN. The most preferred amines for use in the compositions herein are 1-hexylamine, 1-octylamine, 1-decylamine, 1-dodecylamine. Especially desirable for odor characteristics are n-dodecyl-dimethylamine (ARMEEN DM12D) and bishydroxyethyl-coconutalkylamine (ETHOMEEN C/12, BEROL 307) and oleylamine 7 times ethoxylated (BEROL 28), lauryl amido propylamine and coco amido propylamine.

Other suitable amines include tertiary amines having the formula



wherein R_1 is C_4 - C_{10} , preferably C_8 - C_{10} alkyl; n is 2-4, preferably n is 3; R_2 is C_1 - C_4 or



$(\text{CH}_2 - \text{CH} - \text{O})_x \text{H}$, whereby x is 1-5, R_3 is H or C_1 - C_2 alkyl.

When included therein, the laundry detergent compositions of the present invention typically comprise nonionic surfactants in the weight ratio of anionic surfactant to nonionic surfactant from 6:1 to 1:3, preferably from 5:1 to 2:1.

When included therein, the laundry detergent compositions of the present invention typically comprise nonionic surfactants in the weight ratio of anionic surfactant to nonionic surfactant from 6:1 to 1:3, preferably from 5:1 to 2:1.

The compositions according to the present invention may further comprise a builder system. Any conventional builder system is suitable for use herein including aluminosilicate materials, silicates, polycarboxylates and fatty acids, materials such as ethylenediamine tetra-acetate, metal ion sequestrants such as aminopoly-phosphonates, particularly ethylenediamine tetramethylene phosphonic acid and diethylene triamine pentamethylene phosphonic acid. Though less preferred for obvious environmental reasons, phosphate builders can also be used herein.

Suitable polycarboxylates builders for use herein include citric acid, preferably in the form of a water-soluble salt, derivatives of succinic acid of the formula $R-CH(COOH)CH_2(COOH)$ wherein R is C10-20 alkyl or alkenyl, preferably C12-16, or wherein R can be substituted with hydroxyl, sulfo sulfoxyl or sulfone substituents. Specific examples include lauryl succinate, myristyl succinate, palmityl succinate 2-dodecenylsuccinate, 2-tetradecenyl succinate. Succinate builders are preferably used in the form of their water-soluble salts, including sodium, potassium, ammonium and alkanolammonium salts.

Other suitable polycarboxylates are oxodisuccinates and mixtures of tartrate monosuccinic and tartrate disuccinic acid such as described in US 4,663,071.

Especially for the liquid execution herein, suitable fatty acid builders for use herein are saturated or unsaturated C10-18 fatty acids, as well as the corresponding soaps. Preferred saturated species have from 12 to 16 carbon atoms

in the alkyl chain. The preferred unsaturated fatty acid is oleic acid. Other preferred builder system for liquid compositions is based on dodecenyl succinic acid and citric acid.

Detergency builders are normally included in amounts of from 3% to 50% by weight of the composition preferably from 5% to 30% and most usually from 5% to 25% by weight.

Optional detergent ingredients :

Preferred detergent compositions of the present invention may further comprise one or more enzymes which provide cleaning performance and/or fabric care benefits. Said enzymes include enzymes selected from cellulases, hemicellulases, peroxidases, proteases, gluco-amylases, amylases, lipases, cutinases, pectinases, xylanases, reductases, oxidases, phenoloxidasases, lipoxygenases, ligninases, pullulanases, tannases, pentosanases, malanases, β -glucanases, arabinosidasases or mixtures thereof.

A preferred combination is a detergent composition having a cocktail of conventional applicable enzymes like protease, amylase, lipase, cutinase and/or cellulase in conjunction with the lipolytic enzyme variant D96L at a level of from 50 LU to 8500 LU per liter wash solution.

The cellulases usable in the present invention include both bacterial or fungal cellulase. Preferably, they will have a pH optimum of between 5 and 9.5. Suitable cellulases are disclosed in U.S. Patent 4,435,307, Barbesgoard et al, which discloses fungal cellulase produced from *Humicola insolens*. Suitable cellulases are also disclosed in GB-A-2.075.028; GB-A-2.095.275 and DE-OS-2.247.832.

Examples of such cellulases are cellulases produced by a strain of *Humicola insolens* (*Humicola grisea* var. *thermoidea*), particularly the *Humicola* strain DSM 1800.

Other suitable cellulases are cellulases originated from *Humicola insolens* having a molecular weight of about 50KDa, an isoelectric point of 5.5 and containing 415 amino acids. Especially suitable cellulases are the cellulases having color care benefits. Examples of such cellulases are cellulases described in European patent application No. 91202879.2, filed November 6, 1991 (Novo).

Peroxidase enzymes are used in combination with oxygen sources, e.g. percarbonate, perborate, persulfate, hydrogen peroxide, etc. They are used for "solution bleaching", i.e. to prevent transfer of dyes or pigments removed from substrates during wash operations to other substrates in the wash solution. Peroxidase enzymes are known in the art, and include, for example, horseradish peroxidase, ligninase, and haloperoxidase such as chloro- and bromo-peroxidase. Peroxidase-containing detergent compositions are disclosed, for example, in PCT International Application WO 89/099813 and in European Patent application EP No. 91202882.6, filed on November 6, 1991.

Said cellulases and/or peroxidases are normally incorporated in the detergent composition at levels from 0.0001% to 2% of active enzyme by weight of the detergent composition.

Preferred commercially available protease enzymes include those sold under the tradenames Alcalase, Savinase, Primase, Durazym, and Esperase by Novo Nordisk A/S (Denmark), those sold under the tradename Maxatase, Maxacal and Maxapem by Gist-Brocades, those sold by Genencor International, and those sold under the tradename Opticlean and Optimase by Solvay Enzymes. Also proteases described in our co-pending application USSN 08/136,797 can be included in the detergent composition of the invention. Protease enzyme may be incorporated into the compositions in accordance with the invention at a level of from 0.0001% to 2% active enzyme by weight of the composition.

A preferred protease herein referred to as "Protease D" is a carbonyl hydrolase variant having an amino acid sequence not found in nature, which is derived from a precursor carbonyl hydrolase by substituting a different amino acid for the amino acid residue at a position in said carbonyl hydrolase equivalent to position +76, preferably also in combination with one or more amino acid residue positions equivalent to those selected from the group consisting of +99, +101, +103, +104, +107, +123, +27, +105, +109, +126, +128, +135, +156, +166, +195, +197, +204, +206, +210, +216, +217, +218, +222, +260, +265, and/or +274 according to the numbering of *Bacillus amyloliquefaciens* subtilisin, as described in the concurrently filed patent application of A. Baeck et al. entitled "Protease-Containing Cleaning Compositions" having U.S. Serial No. 08/322,676, filed October 13, 1994, which is incorporated herein by reference in its entirety.

Highly preferred enzymes that can be included in the detergent compositions of the present invention include lipases. It has been found that the cleaning performance on greasy soils is synergistically improved by using lipases. Suitable lipase enzymes include those produced by microorganisms of the *Pseudomonas* group, such as *Pseudomonas stutzeri* ATCC 19.154, as disclosed in British Patent 1,372,034. Suitable lipases include those which show a positive immunological cross-reaction with the antibody of the lipase, produced by the microorganism *Pseudomonas fluorescens* IAM 1057. This lipase is available from Amano Pharmaceutical Co. Ltd., Nagoya, Japan, under the trade name Lipase P "Amano," hereinafter referred to as "Amano-P". Further suitable lipases are lipases such as M1 Lipase^R and Lipomax^R (Gist-Brocades). Highly preferred lipases are the D96L lipolytic enzyme variant of the native lipase derived from *Humicola lanuginosa* as described in US Serial No. 08/341,826. Preferably the *Humicola lanuginosa* strain DSM

4106 is used. This enzyme is incorporated into the composition in accordance with the invention at a level of from 50 LU to 8500 LU per liter wash solution. Preferably the variant D96L is present at a level of from 100 LU to 7500 LU per liter of wash solution. More preferably at a level of from 150 LU to 5000 LU per liter of wash solution.

By D96L lipolytic enzyme variant is meant the lipase variant as described in patent application WO 92/05249 viz. wherein the native lipase ex *Humicola lanuginosa* aspartic acid (D) residue at position 96 is changed to Leucine (L). According to this nomenclature said substitution of aspartic acid to Leucine in position 96 is shown as : D96L.

Also suitable are cutinases [EC 3.1.1.50] which can be considered as a special kind of lipase, namely lipases which do not require interfacial activation. Addition of cutinases to detergent compositions have been described in e.g. WO-A-88/09367 (Genencor).

The lipases and/or cutinases are normally incorporated in the detergent composition at levels from 0.0001% to 2% of active enzyme by weight of the detergent composition.

Amylases (& and/or β) can be included for removal of carbohydrate-based stains. Suitable amylases are Termamyl^R (Novo Nordisk), Fungamyl^R and BAN^R (Novo Nordisk).

The above-mentioned enzymes may be of any suitable origin, such as vegetable, animal, bacterial, fungal and yeast origin.

Said enzymes are normally incorporated in the detergent composition at levels from 0.0001% to 2% of active enzyme by weight of the detergent composition.

Other suitable detergent ingredients that can be added are enzyme oxidation scavengers which are described in Copending European Patent application 92870018.6 filed on January 31, 1992. Examples of such enzyme oxidation scavengers are ethoxylated tetraethylene polyamines.

Other components used in detergent compositions may be employed, such as soil-suspending agents, soil-release polymers, abrasives, bactericides, tarnish inhibitors, coloring agents, foam control agents, corrosion inhibitors and perfumes.

Preferably, the liquid compositions according to the present invention are in "concentrated form"; in such case, the liquid detergent compositions according to the present invention will contain a lower amount of water, compared to conventional liquid detergents. The level of water is less than 50%, preferably less than 30% by weight of the detergent compositions.

Said concentrated products provide advantages to the consumer, who has a product which can be used in lower amounts and to the producer, who has lower shipping costs.

The liquid compositions are especially effective when applied directly to soils and stains in a pretreatment step.

The detergent compositions of the present invention can also be used as detergent additive products. Such additive products are intended to supplement or boost the performance of conventional detergent compositions.

The detergent compositions according to the present invention include compositions which are to be used for cleaning of substrates, such as fabrics, fibers, hard surfaces, skin etc., for example hard surface cleaning compositions (with or without abrasives), laundry detergent compositions, automatic and non-automatic dishwashing compositions.

The following examples are meant to exemplify compositions of the present inventions, but are not necessarily meant to limit the scope of the invention.

EXAMPLE I

The following liquid detergent compositions are made :

% by weight of the detergent compositions

	A	B	C	D
C ₁₂ -C ₁₅ alkyl ethoxylated sulfate	2	8	11	5
C ₈ -C ₁₀ hydroxyethyl dimethyl quat	2	2	2	1
C ₁₂ -C ₁₄ alkyldimethyl amine oxide	-	-	-	2
C ₁₂ -C ₁₅ alkyl sulfate	17	12	7	8
C ₁₂ -C ₁₄ N-methyl glucamide	5	4	4	3
C ₁₂ -C ₁₄ fatty alcohol ethoxylate	6	1	1	1
C ₁₂ -C ₁₈ fatty acid	11	4	4	3
Citric acid anhydrous	1	3	3	2
Diethylene triamine penta methylene phosponic acid	1	1	1	0.5
Monoethanolamine	8	5	5	2
Sodium hydroxide	1	2.5	1	1.5
Propanediol	14.5	13.1	10.0	8
Ethanol	1.8	4.7	5.4	1
Amylase (300KNU/g)	0.1	0.1	0.1	0.1
Lipase D96/L (100KNU/g)	0.15	0.15	0.15	0.15
Protease (35g/l)	0.5	0.5	0.5	0.5)
Endo-A (5000 CEVU/g)	0.05	0.05	0.05	0.5
Carezyme (5000 CEVU/g)	0.09	0.09	0.09	0.9
Terephthalate-based polymer	0.5	-	0.3	0.3
Boric acid	2.4	2.8	2.8	2.4
Sodium xylene sulfonate	-	3	-	-
DC 3225C	0.04	0.04	0.03	0.03
2-butyl-octanol	1	1	1	1
Branched silicone	0.3	0.3	0.3	0.3


Water & minors

Up to 100%

The above liquid detergent compositions (A-D) are found to be very efficient in the removal of greasy/oily soils under various usage conditions while having a controlled suds profile.


What is claimed is:

1. A liquid laundry detergent composition comprising a Nil-Las surfactant system said surfactant system comprising
 - i) anionic surfactants selected from the group of alkyl alkoxy sulfates and alkyl sulfates,
 - ii) a quaternary ammonium surfactant having the formula



Formula I

whereby R1 is a short chainlength alkyl (C6-C10) or alkylamidoalkyl of the formula (II) :



Formula II


y is 2-4, preferably 3.

whereby R2 is H or a C1-C3 alkyl,

whereby x is 0-4, preferably 0-2, most preferably 0,

whereby R3, R4 and R5 are either the same or different and can be either a short chain alkyl (C1-C3) or alkoxyalkyl of the formula III,

whereby X⁻ is a counterion, preferably a halide, e.g. chloride or methylsulfate.



Formula III

R6 is C₁-C₄ and z is 1 or 2, characterized in that the weight ratio of the quaternary ammonium compound to the alkyl alkoxy sulfates and alkyl sulfates is from 1:3 to 1:30, preferably from 1:3 to 1:15, most preferred from 1:5 to 1:10.

2. A liquid detergent composition according to claim 1 having a quaternary ammonium surfactant as defined in claim 1 wherein

R₁ is C₈, C₁₀ or mixtures thereof, x=0,
R₃, R₄ = CH₃ and R₅ = CH₂CH₂OH.

3. A liquid detergent composition according to Claim 1-2 wherein the anionic surfactants are selected from the alkyl ethoxylated sulfate and C₁₂-C₁₅ alkyl sulfate.

4. A liquid detergent composition according to claims 1-3 further comprising a lipolytic enzyme.

5. A liquid detergent composition according to claim 4 wherein said lipolytic enzyme is D96L lipolytic enzyme.

6. A liquid detergent composition according to Claims 1-5 further comprising a nonionic surfactant selected from the polyhydroxy fatty acid amides and/or amine oxides.

7. A liquid detergent composition according to claim 6 wherein said amine oxide is C₁₂-C₁₄ alkyl dimethyl amine oxide.

8. A liquid detergent composition according to Claims 1-7 further comprising other surfactants, builders, enzymes and other conventional detergent ingredients.

9. Use of a liquid detergent composition according to Claims 1-8 for pretreatment of fabrics.

INTERNATIONAL SEARCH REPORT

International application No
PCT/US 95/12470

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C11D1/65 C11D1/86

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C11D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US,A,4 235 759 (JOHNA NOBUO ET AL) 25 November 1980 see claims; example 1 ---	1-3,8
X	EP,A,0 095 205 (PROCTER & GAMBLE) 30 November 1983 see page 13; claims; examples ---	1,3,8
X	EP,A,0 000 224 (PROCTER & GAMBLE) 10 January 1979 cited in the application see page 7; claims; examples 6,12-14 ---	1,2
X	FR,A,2 329 746 (PROCTER & GAMBLE) 27 May 1977 see claims; examples ---	1,6,7
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *&* document member of the same patent family

Date of the actual completion of the international search

31 May 1996

Date of mailing of the international search report

25-06-1996

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Pfannenstein, H

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 95/12470

C(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	W0,A,92 05249 (NOVO NORDISK AS) 2 April 1992 cited in the application see example 20 ---	1,4,5
E	W0,A,95 29218 (PROCTER & GAMBLE) 2 November 1995 see page 6; claims; examples -----	1,3,4,6, 8

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 95/12470

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-4235759	25-11-80	JP-C- 1543570	15-02-90
		JP-A- 54159416	17-12-79
		JP-B- 63037158	22-07-88

EP-A-0095205	30-11-83	AU-B- 559791	19-03-87
		AU-B- 1488883	01-12-83
		CA-A- 1208519	29-07-86
		JP-C- 1785137	31-08-93
		JP-B- 4077038	07-12-92
		JP-A- 59025894	09-02-84
		US-A- 4561998	31-12-85

EP-A-0000224	10-01-79	AT-B- 400244	27-11-95
		AU-B- 525063	21-10-82
		AU-B- 3758778	03-01-80
		BE-T- 7	07-12-79
		BE-T- 8	07-12-79
		CA-A- 1109759	29-09-81
		CA-A- 1129751	17-08-82
		CH-A- 643881	29-06-84
		DE-A- 2857161	06-03-80
		DE-A- 2857163	06-03-80
		EP-A- 0000225	10-01-79
		FR-A,B 2423534	16-11-79
		FR-A,B 2423537	16-11-79
		GB-A,B 2040986	03-09-80
		GB-A,B 2040987	03-09-80
		JP-A- 54039416	26-03-79
		JP-A- 54039417	26-03-79
		NL-T- 7815014	31-10-79
		NL-T- 7815015	31-10-79
		SE-B- 446460	15-09-86
		SE-A- 7907257	31-08-79
		US-A- 4321165	23-03-82
		US-A- 4333862	08-06-82

FR-A-2329746	27-05-77	BE-A- 847843	29-04-77
		DE-A- 2648304	05-05-77
		JP-A- 52086410	18-07-77
		NL-A- 7612025	03-05-77

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 95/12470

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
W0-A-9205249	02-04-92	AU-B- 657278	09-03-95
		AU-B- 8617291	15-04-92
		CA-A- 2092615	14-03-92
		EP-A- 0548228	30-06-93
		JP-T- 6501153	10-02-94

W0-A-9529218	02-11-95	US-A- 5466394	14-11-95
